



Exploring the Significance and Medicinal Potential of *Rubus fraxinifolius*: A Review of Ragimot Wildberry

Rubus fraxinifolius Poir.

Muhamad Hanif Rawi¹

¹ Universiti Malaysia Sabah [Please enter the full address including contact form \(email, phone\).](#)

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract [The popular name varies in different regions or countries. Therefore, please, throughout the text, replace the popular name with the scientific name.](#)

The rainforests of Sabah, Malaysia, are home to a diverse array of plant life, including the Ragimot berry (*Rubus fraxinifolius*). This ~~berry~~ ^{plant}, belonging to the *Rubus* genus, is nutritionally rich and holds important cultural and medicinal ~~importance~~ ^{importance}. Traditionally, *Rubus* species have been known for their various medicinal properties, such as anti-inflammatory, antioxidant, anti-tumor, and neuroprotective effects. The ~~Ragimot~~ ^{*Rubus fraxinifolius*} berry contains high levels of sugar, vitamin C, and iron, contributing to its health-promoting qualities. It also exhibits strong antioxidant activity and potential as an anti-acetylcholinesterase and antibacterial agent, suggesting its use in managing neurodegenerative diseases and as a natural antimicrobial substance. Efforts are currently underway to domesticate Ragimot through research focused on optimizing cultivation practices. Scientists are studying the best timing and dosage for fertilizer application and exploring methods for large-scale propagation through in vitro germination and plant regeneration strategies. The goal of this research is to conserve this unique plant species endemic to Borneo and provide **economic benefits** to local communities. To summarize, ~~Ragimot~~ ^{*Rubus fraxinifolius*} is a valuable plant found in Sabah's rainforests. It is rich in nutrients and has functional properties that make it promising for future use. More research is needed to fully understand its health benefits and promote responsible use. ~~Ragimot~~ ^{*Rubus fraxinifolius*} serves as a reminder of the importance of protecting biodiversity while utilizing it for a healthier and more sustainable world.

Introduction

Rubus fraxinifolius, popularly known as ragimot

In the lush rainforests of Sabah, Malaysia, thrives a hidden gem – the ragimot, a wild berry with a rich history and promising potential. This ~~chapter~~ ^{work or study} aims to explore the nutritional composition and functional properties of ~~ragimot~~ ^{*Rubus fraxinifolius* Poir.}, shedding light on its significance in the domains of food and medicine. As global interest in natural and sustainable resources continues to rise, understanding the attributes of ragimot becomes crucial for both local communities and the broader scientific community. Known as ~~rogimot~~ ^{*Rubus fraxinifolius*, popularly known as} (pronounced differently due to Dusun and Kadazandusun dialect influences) in Sabah, ~~this~~ ^{its} berry is favoured by climbers despite its somewhat tasteless reputation. It belongs to the *Rubus* subgenus *Ideobatus*, and its fruits are sold in markets and made into jams, particularly in Java (Lamb, 2019). More than eight species of *Rubus* are found above 1200 m on Mount Kinabalu (Corner & Beaman, 1996), including *Rubus fraxinifolius*. Local variations of ~~ragimot~~ ^{*Rubus fraxinifolius*} come in two forms: one is red and oval-shaped, and the second is red and round-shaped. However, for this ~~chapter~~ ^{work}, the focus is on the former, *Rubus fraxinifolius*. The *Rubus* genus, encompassing 740 species ~~within Rosaceae~~ ^{of the family Rosaceae}, has a remarkable presence on six continents, thriving in diverse habitats. Within the Malesia Region, 46 *Rubus* species were discovered. One of these is *R. fraxinifolius*, a wild raspberry variant. It is found in Sabah's mountainous forests, particularly in the regions around Mount Kinabalu (Shamsudin et al., 2019). Interestingly, the traditional medicinal application of *Rubus* in Southeast Asia closely mirrors its use in Europe. Numerous *Rubus* species have been recognized for producing diverse compounds that exhibit pharmacological properties, such as antidiabetic, antibacterial, anti-inflammatory, antioxidant, antidiarrheal, anti-tumor, neuroprotective, and wound-healing effects. Efforts to domesticate the plants are underway by small farmers in Kundasang, Sabah. While the Cibodas Botanical Garden in Jakarta, Indonesia, a hub for ex situ conservation, has amassed eleven *Rubus* species, including *R. fraxinifolius* (Dewi et al., 2019). Based on data collected from the Cibodas Botanical Garden, *arben* (the common name known in Indonesia) is available throughout the year, with its peak season occurring from January to March (Surya & Rahman, 2012). The flowering and fruiting patterns of the plants correlate positively with environmental factors, such as monthly relative humidity. This observation aligns with the fluctuating availability of ragimot in Kundasang, Sabah, depending on the wet seasons in the area. In the realm of the animal kingdom, the babirusa, a swine-related mammal, feeds on the fruits of *R. fraxinifolius*. *Babyrousa quadricornua*, knowed in India as babirusa *R. fraxinifolius* ~~ragimot~~ found in Maluku, Indonesia (Macdonald et al., 2023). The ~~ragimot~~ ^{*R. fraxinifolius*} berry (~~*Rubus* spp.~~ ^{*Rubus* spp.}) itself is an erect shrub, standing 2-3 meters tall and adorned with up to 6 mm prickles on its stems. Its pinnate leaves hold 4 pairs of opposite leaflets and a terminal one. These elliptic leaflets, spanning 2-9 x 1.4 cm, sport serrated edges and are sparsely covered in hair, with 7-10(-15) vein pairs (Lamb, 2019). Inflorescence panicles, measuring 6-20 cm long, bear grayish yellow-green flowers with bulging forms (Normasiwi et al., 2021). The small drupes, ranging in color from orange to red, cluster on a central stalk and can measure up to 2.5 x 1.5 cm, each containing a tiny seed. The ~~ragimot~~ ^{*R. fraxinifolius*} range extends from Taiwan (Huang & Hu, 2009), the Philippines, and Borneo to Sulawesi, Lawu Mountain in East Java (Hidayah & Roziaty, 2022), New Guinea, the Solomon Islands, and even Le Pouce Mountain Reserve in Mauritius, albeit in low abundance (Bissessur et al., 2023).

What is the difference between the *R. fraxinifolius* specie of the *Rubus fraxinifoliolus* Hayata?

Nutritional composition Please avoid repetitive information, specify the species being studied, highlight the data that differentiate this species from the others. Do this throughout the text.

Rubus fraxinifolius

Ragimot is a species of wild raspberry found in the mountainous forests (Desmiaty & Elya, 2021). The berry is valued for its taste, medicinal properties, and nutritional composition (Jansen-Alves et al., 2021). Ragimot is rich in phytochemicals, which are bioactive compounds that contribute to its health-promoting properties. The fruit of Ragimot has been found to have a high content of sugar, vitamin C, and iron (Surya et al., 2018). Specifically, it has been reported to contain 5.05 g of sugar per 100 g of fruit, which is higher than other species (*R. rosifolius*, *R. chrysophyllus*, *R. pyrofolius*, and *R. idaeus*) of wild *Rubus* (Surya et al., 2018). In terms of vitamin C content, Ragimot has been found to have the highest amount (83.65 mg/100 g) during the ripening fruit stage II (Surya et al., 2018). Additionally, it has been reported to have a considerable content of iron (Surya et al., 2018). In terms of phytochemical composition, Ragimot has been found to

contain total phenolics, flavonoids, and carotenoids (Bakar et al., 2016). These phytochemicals contribute to the antioxidant activity of the fruit. Antioxidants play a crucial role in protecting the body against oxidative stress and reducing the risk of chronic diseases (Bakar et al., 2016). The antioxidant activity of Ragimot has been evaluated using various assays, including the 1,2-diphenyl-2-picrylhydrazyl (DPPH), iron-reducing antioxidant power (FRAP), and 2,2-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) assays (Bakar et al., 2016). These assays measure the ability of the fruit extract to scavenge free radicals and inhibit oxidative damage. Furthermore, Ragimot has been found to have anti-acetylcholinesterase and antibacterial activities (Bakar et al., 2016). These activities suggest potential therapeutic applications of the fruit in the treatment of neurodegenerative diseases and as a natural antimicrobial agent. Extracts from the leaves contain alkaloids, flavonoids, hydrolyzed tannins, saponins, and triterpenoids (Sulistyowati et al., 2023). Based on the data, the old leaves yield a significantly higher amount of phytochemicals than the younger leaves. These natural antioxidants scavenge free radicals, helping to combat oxidative stress when used as an antiaging agent. As reported by Dewi et al. (2019) in *Rubus fraxinifolius* Hayata, fresh leaves had a higher content of phenolics and flavonoids and antioxidant capacity than dry leaves, but the α -glucosidase inhibitory activity was higher in air-dried leaves.

Rubus fraxinifolius Hayata. Please, see The Word Flora Online data.

Functional properties

Rubus fraxinifolius ?? This information has been mentioned before. ???
Ragimot stands out as an extraordinary fruit that has garnered substantial interest due to its exceptional nutritional composition and potential health advantages. This indigenous Borneo plant, cherished for its year-round harvestability, has emerged as a critical source of income for local communities in Sabah. The fruit's remarkable attributes, such as its elevated sugar, vitamin C, and iron content, not only contribute to its nutritional value but also position it as an invaluable dietary addition (Wardah et al., 2022). *Rubus fraxinifolius* holds deep ethnomedicinal significance, as indigenous communities, such as the Kankanaeys and Kalanguyas in the Philippines, have incorporated the berry into their dietary practices and traditional treatments (Bersamin et al., 2021; Galvez, 2015; Gomez Jr et al., 2023). For instance, the Kalanguyas utilize crushed leaves for managing sore eyes and wounds, while root, leaf, and trunk decoctions are employed to address issues like diarrhea and urinary tract infections (Gomez Jr et al., 2023). This multifaceted plant is being explored from various angles, including its stems, leaves, and fruits, revealing anthocyanin, ellagitannin, proanthocyanin (Desmiaty et al., 2019), and potent triterpenoids with inhibitory actions. These triterpenoids contribute to

the plant's elastase inhibitory activity, expanding our comprehension of its bioactive potential (Desmiaty et al., 2020).

Beyond its nutritional richness, *R. fraxinifolius* offers an array of functional properties that promote well-being. Both its leaves and fruits exhibit remarkable radical scavenging activity, showcasing its potential as a natural reservoir of antioxidants (Desmiaty et al., 2018). This prowess in scavenging free radicals is attributed to the abundant polyphenols and flavonoids present in Ragimot. These compounds possess potent antioxidant capabilities that contribute to the fruit's potential role in traditional medicines and anti-aging cosmetics. Notably, recent studies (Sulistiyowati et al., 2023) emphasize the efficacy of old leaves in displaying superior antioxidant and elastase inhibition activity when compared to their younger counterparts. These findings suggest the feasibility of harnessing *R. fraxinifolius* leaves as an active ingredient in anti-aging cosmetic formulations. The fruit's potent ability to scavenge radicals has been demonstrated through the FRAP and DPPH methods, underscoring its antioxidant potential (Desmiaty & Elya, 2021). The rich reservoir of polyphenols in the fruit contributes significantly to this antioxidant capacity, positioning *R. fraxinifolius* as a superior candidate in comparison to other *Rubus* species (Bakar et al., 2016). *R. fraxinifolius* stands out for its potential as an anti-aging agent, demonstrating anti-elastase and anti-tyrosinase activities that are pivotal in thwarting skin aging (Desmiaty et al., 2020). These attributes render it a promising contender for incorporation into cosmetics and skincare formulations aimed at preserving youthful skin. Furthermore, the species showcases nutritional prowess, boasting elevated sugar, vitamin C, and iron content, setting it apart from other *Rubus* counterparts (Surya et al., 2018). Amid its diverse functional properties, *R. fraxinifolius* has also been investigated for its antibacterial potential. While some *Rubus* species exhibit varying inhibitory effects on different bacterial strains (Bakar et al., 2016), as shown by Galvez et al. (2016), ragimot showed antibacterial potential against *Bacillus cereus*. *R. fraxinifolius*'s antibacterial properties underscore its versatility and multifaceted contributions. In the realm of scientific inquiry, *R. fraxinifolius* has emerged as a captivating subject, revealing a wealth of functional attributes that span from antioxidant and anti-aging potential to antibacterial efficacy. These revelations herald a new era of exploration into the possibilities that this remarkable plant holds across domains as diverse as medicine, cosmetics, and dietary enhancement. As we unravel its mysteries, *R. fraxinifolius* cements its place as an invaluable botanical treasure.

Please provide more data confirming this statement, with the respective references.

Efforts on cultivation

Addressing the urgent need to conserve a rapidly declining Borneo-endemic plant species, *R. fraxinifolius* Ragimot, has prompted extensive research efforts (Bissessur et al., 2023). In pursuit of this mission, recent studies have investigated various aspects of Ragimot cultivation. Notably, a groundbreaking approach centered on optimizing fertilizer application time and dosage has been undertaken to domesticate and cultivate *R. fraxinifolius* Ragimot as a highly promising fruit crop (Noviady et al., 2022). This endeavor builds upon prior research, which had already identified compost as the optimal planting medium, demonstrating its efficacy across multiple parameters (Surya, 2012).

In the context of *Rubus rosifolius* Sm. and *R. fraxinifolius* Poir., their growth responses to combinations of fertilizer doses and application timings have been scrutinized across several research articles. Diving into this field, Ismaini et al. (2017) investigated micropropagation methods for *Rubus chrysophyllus* and *R. fraxinifolius*. The study revealed distinct

responses to a medium containing 10 mg/L of IBA, with *R. fraxinifolius* displaying superior results in terms of root length and numbers compared to *Rubus chrysophyllus*. Building on this foundation, Noviady et al. (2022) delved into the growth and development responses of *Rubus rosifolius* Sm. and *R. fraxinifolius* Poir. to the interplay of fertilizer doses and application timings. Strikingly, their findings showcased the prominence of combinations such as *R. rosifolius* + K3 and *R. fraxinifolius* + K4, leading to elevated fruit yield in terms of number, size, and weight per plant.

These ~~comprehensive~~ ^{important} investigations underscore the pivotal role played by fertilizer doses and application timings in shaping the growth and development of *Rubus* species. The revelations offer a valuable blueprint for optimizing productivity and profitability in fruit crop cultivation, particularly focusing on *Rubus rosifolius* Sm. and *R. fraxinifolius* Poir. (Ismaini et al., 2017; Guo et al., 2021; Noviady et al., 2022).

Given the imperative for large-scale plantlet production for commercial purposes, there's a pressing demand for efficient and cost-effective methods. Presently, the propagation of *R. fraxinifolius* predominantly relies on seedlings growing within the garden. However, this conventional approach is time-intensive, as observed by Surya (2012), with seedling heights ranging between 0.96 – 3.56 cm within 12 weeks after sowing. In contrast, the in vitro germination method, as noted by Surya et al. (2015), offers a swifter alternative. Ismaini and Surya (2017) have also explored diverse plant regeneration avenues, honing in on *R. fraxinifolius* hypocotyls. Their research aimed at optimizing the regeneration process through the assessment of various growth regulators' impact on callus dimensions, leaf numbers, plantlet traits, and root counts.

Encouragingly, the results underscored the efficacy of medium 3, comprising 0.5 mg/L GA3, 2.5 mg/L 2-iP, and 0.1 mg/L ^{gibberellic acid} ^{2-iso-pentenyl adenine (2iP)} ~~naphthalene acetic acid (NAA)~~ NAA, in driving the development of *R. fraxinifolius* explants (Ismaini & Surya, 2017). Importantly, the concentrations of 2-iP and NAA were found to wield a significant influence on the regenerative process.

Beyond its medicinal applications, ^{*R. fraxinifolius*} Ragimot holds multifaceted value for local communities in the Barangays, Philippines. This diverse utility spans offerings such as condiments like jam, jellies, candies, juice, wine, as well as its employment in dyes and decorations (Chua-Barcelo, 2014). As research continues to illuminate the potential of this species, its significance transcends boundaries, promising both ecological preservation and economic enrichment.

Conclusion

~~Ragimot, the wild berry of Sabah, stands as a testament to the wealth of natural resources harbored within the world's rainforests.~~ ^{The} ^{of *R. fraxinifolius*} ^{*R. fraxinifolius*} Its nutritional composition, packed with phytochemicals, vitamins, minerals, and dietary fibre, bestows an array of functional properties that can benefit human health. The antioxidant and anti-inflammatory capabilities of Ragimot hold promise in combating various diseases, while its potential to support the immune system and cardiovascular health further solidifies its value as a functional food. Additionally, its role in managing blood sugar levels indicates potential applications in diabetes management. As we embrace a holistic approach to health and nutrition, incorporating traditional and local fruits like Ragimot into our diets may pave the way towards a healthier, sustainable future. However, further research and studies are warranted to unravel the full extent of Ragimot's health benefits, ensuring its responsible use and conservation for the benefit of current and future generations. In conclusion, Ragimot serves as a vibrant reminder of the treasure trove

of natural goodness found within our planet's biodiversity, urging us to preserve these precious resources while harnessing their potential for a healthier and thriving world.

Figures and Tables **Please mention the Figures and Table 1 in the body of the text.**
And, insert the Figures and Table 1 shortly after your first citation.




Figure 1. Nutrient analysis of Ragimot fruits (Surya et al., 2018)

**Figure 1: Nutritional composition of the fruits of *R. fraxinifolius* (Surya et al., 2018).
Photo from <https://tropical.theferns.info/image.php?id=Rubus+fraxinifolius>**



Figure 2. Freshly harvested Ragimot (Sulap Dahai, 2021)



This is a better option.



Figure 3. Ragimot berry may as well resemble raspberry drupes

Table 1: Phytochemicals of *Rubus fraxinifolius* with potential for medical applications

I suggest inserting a column in the middle, ↓ containing examples of possible applications in the medical field.

Table 1. Phytochemical components of ragimot potential for medicinal applications	
Chemical compounds	References
2-propenoic acid; 2-propenyl esters	(Abu Bakar et al., 2016)
2,6-dihydro-xypyridine (6-hydroxy-2(1H)-pyridinone)	
1,1,2-triacetoxyethane	
3-deoxy-d-mannonic lactone furfural	
1,3-butadiene-1-carboxylic acid	
2,4-dihydroxy-2,5-dimethyl-3(2H)-furan-3-one	(Desmiaty et al., 2021)
2,3-O-ethyleneglycol,19-hydroxyurs-12-en-23,28-dioic acid	
2,3-O-propanediol,19-hydroxyurs- 12-en-28-oic acid	

References

- Abu Bakar, M. F., Ismail, N. A., Isha, A., & Mei Ling, A. L. (2016). Phytochemical Composition and Biological Activities

- of Selected Wild Berries (*Rubus moluccanus* L., *R. fraxinifolius* Poir., and *R. alpestris* Blume). *Evidence-Based Complementary and Alternative Medicine*, 2016, 1–10. <https://doi.org/10.1155/2016/2482930>
- Bersamin, A. T., Tayaben, J. L., Balangcod, K. D., Balangcod, A. K. D., Cendana, A. C., Dom-Ogen, E. T., Licnahan, L. O. C., Siadto, B., Wong, F. M., & Balangcod, T. D. (2021). Utilization of plant resources among the Kankanaeys in Kibungan, Benguet Province, Philippines. *Biodiversitas Journal of Biological Diversity*, 22(1), Article 1. <https://doi.org/10.13057/biodiv/d220144>
 - Bissessur, P., Reinegger, R. D., Baider, C., Mamoodie, R., & Florens, F. B. V. (2023). Invasive alien plant control: The priority to save one of the most rapidly declining island-endemic plant species worldwide. *Journal for Nature Conservation*, 73, 126417.
 - Chua-Barcelo, R. (2014). Ethno-botanical survey of edible wild fruits in Benguet, Cordillera administrative region, the Philippines. *Asian Pacific Journal of Tropical Biomedicine*, 4, S525-38. <https://doi.org/10.12980/APJTB.4.201414B36>
 - Corner, E. J. H. & Beaman, J. H. (1996). The plant life of Kinabalu-an introduction. In: Wong, K.M. & Phillips, A. (eds.). Kinabalu Summit of Borneo. Kota Kinabalu, Malaysia. The Sabah Society.
 - Desmiaty, Y., & Elya, B. (2021). Unripe fruit of *R. fraxinifolius* as a potential source of antioxidant and antielastase agent. *International Journal of Applied Pharmaceutics*, 78–81. <https://doi.org/10.22159/ijap.2021.v13s2.15>
 - Desmiaty, Y., Elya, B., Saputri, F. C., Dewi, I. I., & Hanafi, M. (2019). Pengaruh Metode Ekstraksi Terhadap Kandungan Senyawa Polifenol dan Aktivitas Antioksidan pada *R. fraxinifolius*. *Jurnal Ilmu Kefarmasian Indonesia*, 17(2), Article 2. <https://doi.org/10.35814/jifi.v17i2.755>
 - Desmiaty, Y., Elya, B., Saputri, F. C., Hanafi, M., & Prastiwi, R. (2018). Antioxidant Activity of *R. fraxinifolius* Poir. And *Rubus rosifolius* J. Sm. Leaves. *Journal of Young Pharmacists*, 10(2s), S93–S96. <https://doi.org/10.5530/jyp.2018.2s.18>
 - Desmiaty, Y., Hanafi, M., Saputri, F. C., Elya, B., Rifai, E. A., & Syahdi, R. R. (2021). Two triterpenoids from *R. fraxinifolius* leaves and their tyrosinase and elastase inhibitory activities. *Scientific Reports*, 11(1), Article 1. <https://doi.org/10.1038/s41598-021-99970-x>
 - Desmiaty, Y., Saputri, F., Hanafi, M., Prastiwi, R., & Elya, B. (2020). Anti-Elastase, Anti-Tyrosinase and Antioxidant of *R. fraxinifolius* Stem Methanolic Extract. *Pharmacognosy Journal*, 12(2), 271–275. <https://doi.org/10.5530/pj.2020.12.42>
 - Dewi, R. T., Fitria, I., Sundowo, A., Agustian, E., Ismaini, L., Normasiwi, S., Noviady, I., Destri, & Surya, M. I. (2019). Phytochemical Constituent's Comparison Using Various Drying Effects on *R. fraxinifolius* Pour Leaves. *Current Agriculture Research Journal*, 7(3), 310–317. <https://doi.org/10.12944/CARJ.7.3.06>
 - Galvez, M. A. C. (2015). Evaluation of DPPH free radical scavenging activity and phytochemical screening of selected folkloric medicinal plants in Tinoc, Ifugao, Cordillera Administrative Region, Philippines. *International Journal of Scientific and Research Publications*, 5(12), 440-445.
 - Galvez, M. A. C. (2016). Antibacterial activity and phytochemical screening of selected folkloric medicinal plants of Maggok, Hungduan, Ifugao, Cordillera administrative region, Philippines. *Int J Sci Res Publ*, 6(1), 460.
 - Gomez Jr, R. A., Leung, J. M., Paing, J. N., Dom-ogen, E. T., Baniaga, A. V., & Napoleon, R. D. (2023). Plant Diversity in Selected Agro-and Forest Ecosystems in Indigenous Cultural Communities (ICCs) in the Cordillera Region, Northern

- Philippines. In *Plant Diversity in Biocultural Landscapes* (pp. 515–538). Springer.
- Hidayah, A. R., & Roziaty, E. (2022). Keragaman Tanaman Perdu yang Tumbuh di Sepanjang Jalur Pendakian Cemoro Sewu, Magetan. *Prosiding SNPBS (Seminar Nasional Pendidikan Biologi Dan Saintek)*, 413–419.
 - Huang, J.-Y., & Hu, J.-M. (2009). *Revision of Rubus (Rosaceae) in Taiwan* 54(4).
 - Ismaini, L. and Surya, M. (2017). In vitro plant regeneration from hypocotyl of arben (*R. fraxinifolius* Poir.). *Australian Journal of Crop Science*, 11(04), 474-478. <https://doi.org/10.21475/ajcs.17.11.04.359>
 - Ismaini, L., Destri, D., & Surya, M. (2017). Micropropagation of *Rubus chrysophyllus* Reinw. ex Miq. and *R. fraxinifolius* Poir.. *Journal of Tropical Life Science*, 7(1), 72-76. <https://doi.org/10.11594/jtls.07.01.12>
 - Jansen-Alves, C., Rodrigues Pereira, J., Zambiasi, R. C., & Murowaniecki Otero, D. (2022). *Rosaceae rubus rosifolius* Smith: nutritional, bioactive and antioxidant potential of unconventional fruit. *Natural product research*, 1–5. Advance online publication. <https://doi.org/10.1080/14786419.2022.2160979>
 - Lamb, A. (2019). *A guide to wild fruits of Borneo* (1st ed, pp. 226–227). Natural History Publications.
 - Macdonald, A. A., Ziehmer, B., Kitchener, A. C., Gelang, M., Åblad, B., Lintonsson, R., Pücker, K. von, Schaub, S., Kiefer, I., & Schwarz, T. (2023). A Computed Tomographic Study of the Premolar Teeth of *Babyrousa* spp. *Journal of Veterinary Dentistry*, 08987564231166551.
 - Normasiwi, S., Salamah, A., & Surya, M. I. (2021). Morphological characteristics of Indonesian *Rubus* flowers. *Biodiversitas Journal of Biological Diversity*, 22(3), Article 3. <https://doi.org/10.13057/biodiv/d220347>
 - Noviady, I., Junaedi, A., Susanto, S., & Surya, M. (2022). Respons pertumbuhan dan perkembangan *Rubus rosifolius* Sm. dan *R. fraxinifolius* Poir. terhadap kombinasi dosis dan waktu pemberian pupuk. *Buletin Kebun Raya*, 25(2), 76-83. <https://doi.org/10.55981/bkr.2022.798>
 - Sánchez-Hernández, E., Teixeira, A., Pereira, C., Cruz, A., Martín-Gil, J., Oliveira, R., & Martín-Ramos, P. (2023). Chemical constituents and antimicrobial activity of a *Ganoderma lucidum* (Curtis.) P. Karst. aqueous ammonia extract. *Plants*, 12(12), 2271. <https://doi.org/10.3390/plants12122271>
 - Shamsudin, N., Matawali, A., & Jualang Azlan, G. (2019). *Comparison of Antioxidant Activity and Phytochemical Content of Borneo Wild Berry, R. fraxinifolius (Rogimot)*. 6, 36–41.
 - Sulap Dahai [@sulapdahai]. (2021, July 15). [Photograph of the freshly harvest ragimot]. Retrieved from https://www.instagram.com/p/CRWHGxCM8tV/?utm_source=ig_web_copy_link&igshid=MzRIODBiNWFIZA==
 - Sulap Dahai [@sulapdahai]. (2021, November 2). [Photograph of the ragimot berry]. Retrieved from https://www.instagram.com/p/CVxO8TqIzD6/?utm_source=ig_web_copy_link&igshid=MzRIODBiNWFIZA==
 - Sulistyowati, B. E., Iswandana, R., & Nur, S. (2023). Phytocompounds and in vitro antiaging activity of ethanolic extract and fractions of *R. fraxinifolius* Poir. Leaves. *Journal of Pharmacy & Pharmacognosy Research*, 11(4), 595–610.
 - Surya, M. I. (2012). Growth Responses of Arben (*R. fraxinifolius* Poir.) Seedling to Various Planting Media. *Biospecies*, 5, 26–30.
 - Surya, M. I., & Rahman, W. (2012). Flowering and Fruiting Phenology of *Rubus* spp. In Cibodas Botanical Garden, Indoensia. *AGRIVITA, Journal of Agricultural Science*, 34(2), Article 2. <https://doi.org/10.17503/agrivita.v34i2.111>
 - Surya, M. I., Suhartati, S., Ismaini, L., Lusini, Y., Anggraeni, D., Normasiwi, S., Asni, N., & Bakar Sidiq, M. A. (2018). Fruit Nutrients of Five Species of Wild Raspberries (*Rubus* spp.) from Indonesian Mountain's Forests. *Journal of*

Tropical Life Science, 8(1).

- Wardah, Sujarwo, W., Setiawan, M., & Satya, I. A. (2022). Community dependence on biodiversity of food sources around the protected area of Mount Jampang forest as a form of conservation and sustainable development in Garut Regency. *IOP Conference Series: Earth and Environmental Science*, 976(1), 012024. <https://doi.org/10.1088/1755-1315/976/1/012024>

To the Teacher
Muhamad Hanif Rawi
I wish you and all who are by your side
a **good year** 2024,
with **health, peace**
and **many achievements**.